arXiv:1609.08668v1 [math.AP] 27 Sep 2016

Singular elliptic equation involving the GJMS operator on the standard unit sphere.

Mohammed Benalili and Ali Zouaoui

September 29, 2016

Abstract

Given a Riemannian compact manifold (M, g) of dimension $n \geq 5$, we have proven in [1] under some conditions that the equation :

$$P_g(u) = Bu^{2^\sharp - 1} + \frac{A}{u^{2^\sharp + 1}} + \frac{C}{u^p} \tag{1}$$

where P_g is GJMS-operator, $n = \dim(M) > 2k$ ($k \in \mathbb{N}^\star$), A, B and C are smooth positive functions on M, $p > 1$ and $2^\sharp = \frac{2n}{n-2k}$ denotes the critical Sobolev of the embedding $H^2_k(M) \subset L^{2^\sharp}(M)$, admits two distinct positive solutions. The proof of this result is essentially based on the given smooth function $\varphi > 0$ with norm $\|\varphi\|_{P_g} = 1$ fulfilling some conditions (see Theorem 3 in [1]). In this note we construct an example of such function on the unit standard sphere $(\mathbb{S}^n, h)$. Consequently the conditions of the Theorem are improved in the case of $(\mathbb{S}^n, h)$.

1 Construction of the function φ

Inspired by the work of F. Robert. (see [4]) , we construct an example of a smooth function $\varphi > 0$ on the Euclidean sphere $(\mathbb{S}^n, h)$ with norm$\|\varphi\|_{P_h} = 1$. Indeed let $\lambda > 0$ and $x_0 \in \mathbb{S}^n$. To a rotation, we may assume that x_0 is the north pole i.e. $x_0 = (0, ...0, 1)$. We consider the transformation

$$\phi_\lambda : \mathbb{S}^n \to \mathbb{S}^n$$

defined by $\phi_\lambda(x) = \psi_{x_0}^{-1}\left(\lambda^{-1}.\psi_{x_0}(x)\right)$ if $x \neq x_0$ and $\phi_\lambda(x_0) = x_0$ where ψ_{x_0} is the stereographic projection of x_0 given by

$$\psi_{x_0} : (\mathbb{S}^n \setminus \{x_0\}, h) \to (\mathbb{R}^n, \xi),$$

for any $a = (\eta_1, ..., \eta_n, \zeta)$ associates $\psi_{x_0}(a) = \left(\frac{\eta_1}{1-\zeta}, ..., \frac{\eta_n}{1-\zeta}\right)$ and

$$\begin{array}{c} \delta_\lambda : (\mathbb{R}^n, \xi) \to (\mathbb{R}^n, \xi) \\ x \mapsto \delta_\lambda(x) = \frac{1}{\lambda} x \end{array}$$

is the homothetic mapping. h is the canonical metric on $\mathbb{S}^n$ and ξ is the Euclidean one on $\mathbb{R}^n$.

Note that ψ_{x_0} is a conformal, mapping more precisely we have

$$\left(\psi_{x_0}^{-1}\right)^\star h = U^{\frac{4}{n-2k}}.\xi$$

where $U(x) = \left(\frac{1+\|x\|^2}{2}\right)^{k-\frac{n}{2}}$. Hence ϕ_λ is conformal i.e.

$$\phi_\lambda^\star h = u_{x_0,\beta}^{\frac{4}{n-2k}}.h \quad \text{where } \beta = \frac{1+\lambda^2}{\lambda^2-1}$$

and

$$u_{x_0,\beta}(x) = \left(\frac{\sqrt{\beta^2-1}}{\beta - \cos d_h(x, x_0)}\right)^{\frac{n-2k}{2}} \qquad \forall x \in \mathbb{S}^n \text{ with } \beta > 1.$$

In particular we have

$$\int_{\mathbb{S}^n} u_{x_0,\beta}^{2^\sharp} dv_h = \omega_n \tag{2}$$

where $\omega_n > 0$ is the volume of the unit standard sphere $(\mathbb{S}^n, h)$.
By the conformal invariance of the operator P_h on $(\mathbb{S}^n, h)$, we obtain that

$$P_h(u_{x_0,\beta}) = \frac{n-2k}{2} Q_h u_{x_0,\beta}^{2^\sharp - 1} \tag{3}$$

where Q_h denotes the Q-curvature of $(\mathbb{S}^n, h)$ which expresses by the Gover's formula as:

$$Q_h = \frac{2}{n-2k} P_h(1) = \frac{2}{n-2k}(-1)^k \prod_{l=1}^{k} (c_l \; Sc)$$

where $c_l = \frac{(n+2l-2)(n-2l)}{4n(n-1)}$, $Sc = n(n-1)$ (the scalar curvature of $(\mathbb{S}^n, h)$).
So the Q_h is a positive constant.
Multiplying the two sides of (3) by $u_{x_0,\beta}$ and integrating on $\mathbb{S}^n$ we get:

$$\int_{\mathbb{S}^n} u_{x_0,\beta} P_h(u_{x_0,\beta}) dv_h = \frac{n-2k}{2} Q_h \int_{\mathbb{S}^n} u_{x_0,\beta}^{2^\sharp} dv_h.$$

And since

$$\int_{\mathbb{S}^n} u_{x_0,\beta} P_h(u_{x_0,\beta}) dv_h = \|u_{x_0,\beta}\|^2_{P_h}$$

(3) writes

$$\|u_{x_0,\beta}\|^2_{P_h} = \frac{n-2k}{2} Q_h \omega_n.$$

Hence by putting

$$\varphi = \left(\frac{n-2k}{2}\omega_n Q_h\right)^{\frac{-1}{2}} u_{x_0,\beta}$$

we obtain a function satisfying the conditions of Theorem 3 in [1] i.e. $\varphi > 0$ smooth on $(\mathbb{S}^n, h)$ such that $\|\varphi\|_{P_g} = 1$

2 Existence results on the sphere

On the standard unit sphere $(\mathbb{S}^n, h)$, if we take the function φ of Theorem 3 in [1] equals $\left(\frac{n-2k}{2}\omega_n Q_h\right)^{\frac{-1}{2}} u_{x_0,\beta}$, we obtain

Theorem 1 *Let $(\mathbb{S}^n, h)$ be the unit standard unit sphere of dimension $n > 2k$, $k \in \mathbb{N}^\star$. There is a constant $C(n,p,k) > 0$ depending only on n, p, k such that*

$$\frac{1}{2^\sharp}\left(\frac{n-2k}{2}\omega_n Q_h\right)^{\frac{2^\sharp}{2}} \int_{\mathbb{S}^n} \frac{A(x)}{u^{2^\sharp}_{x_0,\beta}} dv_h \leq C(n,p,k)\left(S\max_{x\in\mathbb{S}^n} B(x)\right)^{\frac{2+2^\sharp}{2-2^\sharp}} \tag{4}$$

and

$$\frac{1}{p-1}\left(\frac{n-2k}{2}.\omega_n.Q_h\right)^{\frac{p-1}{2}} \int_{\mathbb{S}^n} \frac{C(x)}{u^{p-1}_{x_0,\beta}} dv_h \leq C(n,p,k)\left(S\max_{x\in\mathbb{S}^n} B(x)\right)^{\frac{p+1}{2-2^\sharp}} \tag{5}$$

where

$$u_{x_0,\beta}(x) = \left(\frac{\sqrt{\beta^2-1}}{\beta - \cos d_h(x,x_0)}\right)^{\frac{n-2k}{2}} \qquad \forall x \in \mathbb{S}^n \text{ and } \beta > 1.$$

then the equation (1) admits a solution of class $C^\infty(\mathbb{S}^n)$. If moreover for any $\varepsilon \in]0, \lambda^\star[$ where $\lambda^\star$ is a positive constant the two following conditions are satisfied

$$\frac{2}{a}\left(\int_{\mathbb{S}^n} \sqrt{A(x)} dv_h\right)^2 \left(\frac{1}{t_0 a_1}\right)^{2^\sharp} > 2^\sharp k \frac{t_0^2}{4n}(2-a)$$

and

$$\left(\frac{2}{a}\right)^{\frac{p-1}{2^\sharp}}\left(\int\limits_{\mathbb{S}^n}\sqrt{C(x)}dv_h\right)^2\left(\frac{1}{t_0a_2}\right)^{p-1}>(p-1)k\frac{t_0^2}{4n}(2-a)$$

where a_1, a_2 *are positive constants,* $2^\sharp=\frac{2n}{n-2k}$ *,* $3<p<2^\sharp+1$*. Then the equation* (1) *admits a second solution.*

Note that since

$$\left(\frac{\beta-1}{\beta+1}\right)^{\frac{n-2k}{4}}\leq u_{x_0,\beta}(x)\leq\left(\frac{\beta+1}{\beta-1}\right)^{\frac{n-2k}{4}}\tag{6}$$

we can improve the conditions (4) and (5) of Theorem 1. Indeed, from (**??**) we deduce that

$$\varphi(x)\geq\left(\frac{\beta-1}{\beta+1}\right)^{\frac{n-2k}{4}}\left(\frac{n-2k}{2}\omega_nQ_h\right)^{-\frac{1}{2}}.$$

Consequently

$$\frac{\|\varphi\|^{2^\sharp}}{2^\sharp}\int_{\mathbb{S}^n}\frac{A(x)}{\varphi^{2^\natural}}dv_h=\frac{1}{2^\sharp}\int_{\mathbb{S}^n}\frac{A(x)}{\varphi^{2^\natural}}dv_h\leq\frac{1}{2^\sharp}\left(\frac{\beta+1}{\beta-1}\right)^{\frac{n}{2}}\left(\frac{n-2k}{2}.\omega_n.Q_h\right)^{\frac{2^\sharp}{2}}\int_{\mathbb{S}^n}A(x)dv_h$$

So, if

$$\frac{1}{2^\sharp}\left(\frac{\beta+1}{\beta-1}\right)^{\frac{n}{2}}\left(\frac{n-2k}{2}.\omega_n.Q_h\right)^{\frac{2^\sharp}{2}}\int_{\mathbb{S}^n}A(x)dv_h\leq C\left(n,p,k\right)\left(S\max_{x\in\mathbb{S}^n}B(x)\right)^{\frac{2+2^\sharp}{2-2^\sharp}}$$

then the condition (4) is fulfilled. Likewise if

$$\frac{1}{p-1}\left(\frac{n-2k}{2}\omega_nQ_h\right)^{\frac{p-1}{2}}\left(\frac{\beta+1}{\beta-1}\right)^{\frac{n(p-1)}{2.2^\sharp}}\int_{\mathbb{S}^n}C(x)dv_h\leq C\left(n,p,k\right)\left(S\max_{x\in M}B(x)\right)^{\frac{p+1}{2-2^\sharp}}$$

the condition (5) is also true and we deduce the following result:

Corollary 2 *Let* $(\mathbb{S}^n,h)$ *be the unit standard unit sphere of dimension* $n>2k$*,* $k\in\mathbb{N}^\star$*. There is a constant* $C(n,p,k)>0$ *depending only on* n,p,k *such that*

$$\frac{1}{2^\sharp}\left(\frac{\beta+1}{\beta-1}\right)^{\frac{n}{2}}\left(\frac{n-2k}{2}.\omega_n.Q_h\right)^{\frac{2^\sharp}{2}}\int_{\mathbb{S}^n}A(x)dv_h\leq C\left(n,p,k\right)\left(S\max_{x\in\mathbb{S}^n}B(x)\right)^{\frac{2+2^\sharp}{2-2^\sharp}}\tag{7}$$

and

$$\frac{1}{p-1}\left(\frac{n-2k}{2}.\omega_n.Q_h\right)^{\frac{p-1}{2}}\left(\frac{\beta+1}{\beta-1}\right)^{\frac{n(p-1)}{2.2^\sharp}}\int_{\mathbb{S}^n}C(x)dv_h\leq C\left(n,p,k\right)\left(S\max_{x\in M}B(x)\right)^{\frac{p+1}{2-2^\sharp}} \tag{8}$$

where $\beta>1$. *Then the equation*(1) *admits a solution of class* $C^\infty(\mathbb{S}^n)$. *If moreover for any* $\varepsilon\in\,]0,\lambda^\star[$, *where* $\lambda^\star$ *is a positive constant, the two following assumptions are satisfied*

$$\frac{2}{a}\left(\int\limits_{\mathbb{S}^n}\sqrt{A(x)}dv_h\right)^2\left(\frac{1}{t_0a_1}\right)^{2^\sharp}>2^\sharp k\frac{t_0^2}{4n}(2-a) \tag{9}$$

and

$$\left(\frac{2}{a}\right)^{\frac{p-1}{2^\sharp}}\left(\int\limits_{\mathbb{S}^n}\sqrt{C(x)}dv_h\right)^2\left(\frac{1}{t_0a_2}\right)^{p-1}>(p-1)k\frac{t_0^2}{4n}(2-a) \tag{10}$$

where a_1,a_2 *are positive constants,* $2^\sharp=\frac{2n}{n-2k}$, $3<p<2^\sharp+1$. *Then the equation* (1) *admits a second solution.*

References

[1] M. Benalili, A. Zouaoui, Singular elliptic equation involving the GJMS operator on compact Riemannian manifold, arxiv:1605 03870v1[math. DG] (2016).

[2] C.R. Graham, R. Jenne, L.J. Masson. G.A.J. Sparling, Conformally invariant powers of the Laplacian. I. Existence. J. London Math. Soc. **46** (1992), 557-565.

[3] E. Hebey, F. Pacard, D. Pollack, A variational Analysis of Einstein-scalar Field Lichnerowicz Equations on compact Riemannian Manifolds, Commun. Math. Phys. 278,(2008), 117-132.

[4] F. Robert, Admissible Q-curvatures under isometries for the conformal GJMS operators, Contemporary Mathematics, Volume in the honor of Jean-Pierre Gossez.